

# Bulletin of the Geological Society of Norfolk

No. 53 (2003)

Published September 2004

---

## CONTENTS

---

	Page
Editorial	1
Lee, J.R., Booth, S.J., Hamblin, R.J.O., Jarrow, A.M., Kessler, H., Moorlock, B.S.P., Morigi, A.N., Palmer, A., Pawley, S.J., Riding, J.B. and Rose, J.	
A new stratigraphy for the glacial deposits around Lowestoft, Great Yarmouth, North Walsham and Cromer, East Anglia, UK.	3

---

The Geological Society of Norfolk exists to promote the study and understanding of geology in East Anglia, and holds meetings throughout the year.



Click here, to order a print copy of the Bulletin.  
Please, specify the issue or year required.

Back copies of the Bulletin cost £3.50 each (members); £10.00 each (non-members).

For membership details consult our web site <http://www.norfolkgeology.co.uk>

---

ISSN0143-9286

**A NEW STRATIGRAPHY FOR THE GLACIAL DEPOSITS AROUND  
LOWESTOFT, GREAT YARMOUTH, NORTH WALSHAM  
AND CROMER, EAST ANGLIA, UK.**

**J.R. Lee<sup>1,2,\*</sup>, S.J. Booth<sup>1</sup>, R.J.O. Hamblin<sup>1,2</sup>, A.M. Jarrow<sup>1</sup>, H. Kessler<sup>1</sup>,  
B.S.P. Moorlock<sup>1,2</sup>, A.N. Morigi<sup>1</sup>, A. Palmer<sup>2</sup>, S.J. Pawley<sup>2</sup>, J.B. Riding<sup>1</sup> and J. Rose<sup>2</sup>**

<sup>1</sup> British Geological Survey,  
Nicker Hill, Keyworth, Nottingham, NG12 5GG, UK.

<sup>2</sup> Department of Geography,  
Royal Holloway, University of London, Egham, Surrey, TW20 0EX, UK.

\*email: jrlee@bgs.ac.uk

**ABSTRACT**

*A new stratigraphical scheme for the glacial deposits around Lowestoft, Great Yarmouth, North Walsham and Cromer (east of Weybourne and Edgefield) is presented, based on a combined research programme by British Geological Survey and the Department of Geography, Royal Holloway, University of London. This stratigraphical scheme is founded on evidence derived from sedimentological descriptions, geological mapping and analytical lithological techniques including clast lithological analysis, derived pre-Quaternary palynomorphs and heavy mineralogy. The previously accepted 'North Sea Drift'/'Lowestoft Formation' scheme is abandoned in favour of four formations that relate to assemblages of till units and associated outwash lithofacies, the mapping of major discontinuities and morpho- and techno-stratigraphical associations. The new scheme consists of the Happisburgh, redefined Lowestoft, Sheringham Cliffs and Briton's Lane formations.*